



The psychological impact of COVID-19 lockdown on the wellbeing, learning efficacy and sleeping behaviors of Romanian medical students

Codrin-Constantin Burlacu¹, Vlad-Florin Chelaru¹,
Tudor Cătălin Drugan², Andrei-Vlad Bădulescu¹

1) Iuliu Hațieganu University of Medicine and Pharmacy Cluj-Napoca, Romania

2) Department of Medical Informatics and Biostatistics, Iuliu Hațieganu University of Medicine and Pharmacy Cluj-Napoca, Romania

Abstract

Background and aims. The COVID-19 lockdowns are reported to have negatively influenced the wellbeing and learning efficacy of students. In this context, we analyzed the psychological impact of the COVID-19 quarantine on healthcare students, a subpopulation experiencing high stress levels.

Methods. Our survey-based, cross-sectional study assessed wellbeing indicators, lifestyle and learning behaviors before and during the quarantine for 388 Romanian healthcare students.

Results. Our findings included the increase in phone and social media use, at the expense of formal and independent study time; deteriorations in mood, self-organization capacity and learning efficacy, and increased procrastination behaviors. Unexpectedly, our study identified an improvement in sleep quality and duration. The increase in social media use was less severe among rural students. We identified correlations between study time, online activities (including social media), wellbeing indicators and procrastination.

Conclusion. Our study draws attention to quarantine-induced deteriorations of wellbeing and learning capacity in an important category of students.

Keywords: COVID -19, lockdown, psychological impact, online learning

Introduction

The COVID-19 pandemic initially described on January 7th 2020 by a few case studies of pneumonia of an unfamiliar etiology [1,2] was declared a public health emergency by the WHO by the end of the month [3].

In response to a rate of transmission far exceeding that of other coronaviruses, quarantine and physical isolation were implemented by multiple countries in an attempt to minimize the spread of COVID-19 [2,4].

In Romania, the first case was registered on February 26, followed by a state of emergency and a 30-day lockdown being instituted on March 14 when pre-university education units and higher education units were closed [5].

While national authorities around the world focused on limiting the number

of total and severe cases, the potential psychological threat of lockdowns was comparatively neglected, resulting in a range of psychological consequences such as heightened stress, depression, anxiety, insomnia, confusion, and anger [6–10].

Evidence from previous outbreaks revealed similar negative effects of quarantine on mental health and psychological wellbeing. For example, during the SARS epidemic, researchers reported stress [11–13], low mood [14], irritability [14], emotional exhaustion [15,16], and insomnia [8,14] among quarantined individuals, while worry, anxiety, and somatization were described as negative effects of the Ebola outbreak [17,18].

Disturbances in individuals' daily routine [6,9], reduced face to face contact [6], a deteriorating sleep

DOI: 10.15386/mpr-2411

Manuscript received: 04.11.2021
Received in revised form: 07.06.2022
Accepted: 15.06.2022

Address for correspondence:
Vlad-Florin Chelaru
vladflorinchelaru@gmail.com

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License <https://creativecommons.org/licenses/by-nc-nd/4.0/>

schedule leading to insomnia [8–10,14], subjective lack of energy due to insufficient physical activity [19,20], and emotional detachment from friends [21] are reported, all corresponding to affected indicators of well-being [10,19,22,23]. Concurrent with this decline of well-being, students reported changes in learning behaviors, and a deteriorating study and work performance [9,21].

According to Satici et al., intolerance to uncertainty is the main predictive factor of mental health issues and indirect determinant of low mental wellbeing [9,24]. Uncertainty is the generic term used to refer to unpredictable events that increase individuals' stress, and in this specific situation it can be caused by not knowing the duration of the pandemic, fear of infection, the unfamiliar challenges of online education and exams, among others [20,25].

Low mood, fear of uncertainty, lack of energy due to insufficient sleep, and sedentarism [26] have been identified as causes of procrastination, itself a way of avoiding stress and anxiety [20,27,28].

Besides the added psychological pressure due to quarantine, medical students are frequently exposed to a high rate of stress, depression, anxiety, insomnia, emotional disturbance and burn-out periods, due to academic pressure and a short time to acquire knowledge [10,29–32].

Consequently, we attempted to study the impact of the COVID-19 quarantine in a population which already experiences higher-than-average levels of mental health issues related to psychological stress.

The two assumptions of this study based on the work of Xiao et al. [33], and Beutel et al. [34], were that: 1) online courses and quarantine conditions had a direct effect on medical students' well-being and sleep quality, and 2) decreased sleep quality and low mood affected learning behaviors, possibly mediated by an increase in procrastination behaviors (time spent on entertainment or social media) and a decrease in learning efficacy, organization and capacity to concentrate.

We conducted a survey-based, analytical, observational, cross-sectional study in order to identify and quantify the impact of COVID-19 quarantine conditions on Romanian medical students' lifestyle, learning efficacy, procrastination and wellbeing. In addition, we analyzed whether the impact of quarantine-induced stress was influenced by two relevant demographic factors, namely gender and place of origin (urban/rural), and to what extent, and finally we tried to identify and quantify correlations between students' lifestyle, learning efficacy, procrastination and wellbeing.

Method

We used Google Forms to create a questionnaire regarding multiple aspects of a student's life. The form consisted of 3 sections, one pertaining to demographic information, one for questions related to students' pre-quarantine lifestyle, and one for questions related to their

lifestyle during the quarantine.

The first section obtained information on the following: age, gender, university, study program, year of study, other activities or work, place of origin (urban/rural), and residence. It should be noted that healthcare education in Romania consists of 6 different study programs, ranging from 3 to 6 years: General Medicine (6 years), Dentistry (6 years), Pharmacy (5 years), Nursing (4 years), Dental technology (3 years), Nutrition and Dietetics (3 years).

The second section consisted of questions regarding multiple topics. The respondents first answered a few questions on time spent on different activities: formal study (i.e. lectures and practical courses), independent study (i.e. any time spent studying outside formal study), and other activities. Secondly, the student was asked to rate the following indicators of wellbeing: mood, self-organizing capacity, and learning efficacy. The third subsection referred to phone use and time spent on the Internet, by category (social media, study, reading, entertainment). Finally, a number of sleep-related questions were included: self-assessed time spent asleep, bedtime hours, a number of sleep quality questions adapted from the Pittsburgh Sleep Quality Index (PSQI questionnaire) [35], and a few questions related to partaking in sleep-perturbing behavior namely having variable meal timing [36,37], drinking coffee or energy drinks less than 2 h before bed [38,39], alcohol [40,41], tobacco [42,43], or other substances use [44], administration of sedatives or other types of medicine [45], or other unspecified behaviors as identified by the students.

The third section repeated the questions of the second part, with a few additional items: an initial question regarding the student's residence during quarantine, a final question on the respondents' perceived change in procrastination time (3 possible answers: 1=decrease, 2=no change, 3=increase during quarantine compared to before quarantine) and a number of questions on self-assessed changes in morale as a result of the quarantine.

The form was disseminated through Facebook groups used by students of Romanian medical universities to collaborate and organize their academic activities and was open from April 12th to 21st; 389 entries were obtained, of which one was removed due to having invalid answers. No incentives were provided for the respondents. To ensure anonymity, the authors responsible for statistical analyses did not have access to the respondents' (optionally submitted) emails.

The PSQI-derived questions were used to compute a pre-quarantine and quarantine PSQI-like score (henceforth sleep quality score or SQS). The SQS score was computed using 12 questions, with higher values indicating a worse sleep quality.

Post-retrieval processing included adjusting the scales of pre-quarantine and during-quarantine questions to ensure the comparability of results.

For each response, a series of “differential” variables were defined, computed with the following formula:

$$D_{Var} = Q_{Var} - N_{Var}$$

As such, lower values of the differential mean that during quarantine, the value for the selected variable was lower. If the student slept for 6-8 h before quarantine, then $N_{Sleep} = 7$ and if the same student slept under 4 h during quarantine, then $Q_{Sleep} = 3$. As such, $D_{Sleep} = 3 - 7 = -4$, the student slept about 4 hours less during quarantine than before quarantine.

Data retrieval was done using Google Forms and Google Sheets (parts of Google Docs suite). Microsoft Excel was used for data conversion, formatting, parsing, and computing of differentials. Statistical analyses and graphing were performed in RStudio, using R for Microsoft Windows, version 4.0.2 [46].

We used the following statistical tests: Kolmogorov-Smirnov for testing distribution normality, Wilcoxon (paired and unpaired) for testing differences of ordinal data

between groups, and McNemar for differences in sleep-perturbing factors during quarantine compared to the time before the quarantine.

We considered statistically significant all $p < 0.05$. Even though our data are generally not normally distributed, we will represent the data in the following format:

$$\text{median (1}^{st} \text{ quartile - 3}^{rd} \text{ quartile) mean} \pm SD$$

Results

Demographic data

388 students answered our questionnaire. Out of them, 307 (79.12%) were females, and 81 (20.88%) were males; 322 (83%) came from urban households, while 66 (17%) came from rural areas. The median age was 21 (1st quartile 20, 3rd quartile 22), mean age 21.09, standard deviation 1.63. More detailed demographic data are provided in table I.

Table I. Detailed demographic data for the survey’s respondents.

| Demographic variable | Value | Count | Percentage |
|----------------------|-------------------------|-------|------------|
| Age | 19 | 21 | 5.41 |
| | 20 | 151 | 38.91 |
| | 21 | 105 | 27.06 |
| | 22 | 50 | 12.89 |
| | 23 | 32 | 8.25 |
| | 24 | 17 | 4.38 |
| | 25 | 7 | 1.8 |
| | 26 | 4 | 1.03 |
| University | 37 | 1 | 0.25 |
| | Cluj-Napoca | 245 | 63.14 |
| | Bucharest | 10 | 2.58 |
| | Iași | 104 | 26.8 |
| | Târgu-Mureș | 10 | 2.58 |
| | Sibiu | 16 | 4.12 |
| | Oradea | 2 | 0.52 |
| Study program | Timișoara | 1 | 0.26 |
| | General medicine | 286 | 73.71 |
| | Dentistry | 47 | 12.11 |
| | Pharmacy | 12 | 3.09 |
| | Nursing | 32 | 8.25 |
| | Dental technology | 1 | 0.26 |
| Study year | Nutrition and dietetics | 10 | 2.58 |
| | 1 | 29 | 7.47 |
| | 2 | 225 | 58 |
| | 3 | 76 | 19.59 |
| | 4 | 32 | 8.25 |
| | 5 | 24 | 6.19 |
| Gender | 6 | 2 | 0.52 |
| | Female | 307 | 79.12 |
| Origin | Male | 81 | 20.88 |
| | Urban | 322 | 83 |
| | Rural | 66 | 17 |

In Romania, some Universities provide their students with accommodation via dorms (137 of respondents, 35.31%); nonetheless, some students prefer to rent or live in an apartment if they study in another city than their hometown (137 of respondents, 35.31%). When the virus began to spread, universities turned to the online to continue their educational process. As such, at the time of our questionnaire, 352 students (90.72%) were staying with their family, while 29 (7.47%) remained in the city of their studies.

Observed behavior before and during the quarantine

In table II we present the median, quartiles, mean and standard deviation of time periods dedicated to different activities during the day, time dedicated to online activities, sleep quantity and quality, and the values of wellbeing indicators. While our data do not generally follow the normal distribution (see below), we nonetheless mentioned the mean and standard deviation, as we considered them to be more illustrative to the reader. Bedtime hour is shown as time, while sleep quality is shown as a score (higher values mean worse sleep). Indicators of wellbeing use a 5-step scale, with higher values meaning more positive views about their represented dimensions. Finally, change in procrastination is a 3-step scale (1=decrease during quarantine, 2=no change, 3=increase during quarantine).

Differences between daily time use, procrastination, sleep quality, and wellbeing indicators during quarantine compared to pre-quarantine conditions

We used the Kolmogorov-Smirnov test to confirm

that the data were not normally distributed (all variables were confirmed to not have a normal distribution, $P < 0.001$), and the Wilcoxon signed-rank test with continuity correction was used to confirm the statistical signification of observed differences in the lifestyle and wellbeing of the students during quarantine, compared to pre-quarantine values.

We have evaluated changes in the following variables during quarantine compared to pre-quarantine conditions: formal study time (mean decrease 1.01h, $p < 0.001$), independent study time (mean decrease 0.28 h, $p = 0.002$), time spent with other daily activities (mean increase 1.91 h, $p < 0.001$), phone use (mean increase 1.23 h, $p < 0.001$).

Regarding time spent online, we found a statistically significant ($p < 0.001$) increase in time spent with social media (mean 0.55 h), formal study (mean 0.59 h), and entertainment (mean 0.81 h), but no statistically significant difference in time spent online for independent study (mean 0.04h) during quarantine compared to pre-quarantine conditions.

Regarding sleep and sleep quality, we found the following statistically significant differences ($p < 0.001$): sleep duration (mean increase 0.64 h), bedtime hour (on average, students went to sleep 25 minutes later during quarantine than before), and sleep quality (decreased score i.e. better sleep quality during quarantine). A graphical representation of this data is available in figure 1.

Finally, we found statistically significant decreases in mood, self-organizing capacity, and learning efficacy ($P < 0.001$), meaning that during quarantine the students reported a lower level of well-being during the lockdown.

This data are represented, using the format mentioned before, in table III.

Table II. Time spent with daily and online activities, sleep quality and duration, and wellbeing indicators.

| Variables | Before quarantine | During quarantine |
|------------------------------|---------------------------------|---------------------------------|
| DAILY ACTIVITIES (h) | | |
| Formal study time | 5 (5-7) 5.87±1.35 | 5 (5-5) 4.86±1.56 |
| Independent study time | 3 (1-3) 2.77±1.69 | 3 (1-3) 2.49±1.73 |
| Other daily activities | 1 (1-3) 2.19±1.49 | 3 (3-5) 4.1±1.75 |
| Phone use | 3 (1.5-3) 2.86±1.61 | 3 (3-5) 4.09±1.92 |
| ONLINE ACTIVITIES (h) | | |
| Social media | 1 (1-2.5) 1.79±1.19 | 2.5 (1-3.5) 2.34±1.38 |
| Formal study | 1 (0.33-1) 1.17±1.13 | 1 (0.33-2.5) 1.76±1.43 |
| Independent study | 1 (1-2.5) 1.76±1.2 | 1 (1-2.5) 1.72±1.3 |
| Entertainment | 2.5 (1-2.5) 2.02±1.25 | 3.5 (2.5-3.5) 2.83±1.36 |
| SLEEP | | |
| Sleep time (h) | 7 (7-7) 6.79±0.97 | 7 (7-9) 7.43±1.28 |
| Sleep quality (SQS value) | 7 (3-13) 8.69±6.93 | 6 (2-11.25) 8.08±8.27 |
| Bedtime hour | 00:30 (23:30-00:30) 00:07±1.06h | 00:30 (23:30-01:30) 00:32±1.17h |
| WELLBEING | | |
| Mood | 4 (3-4) 3.69±0.94 | 3 (2-4) 3.07±1.11 |
| Self-org capacity | 4 (3-4) 3.65±0.85 | 3 (2-4) 2.72±1.15 |
| Learning efficacy | 4 (3-4) 3.55±0.82 | 3 (2-3) 2.6±1.12 |
| Change in procrastination | | 3 (2-3) 2.48±0.7 |




Figure 1. Graphical representation of changes in sleep-related behavior. For the boxplot representing sleep quality scores (lower values meaning better sleep), the middle line represents the median; the boxplot margins mean 1st and 3rd quartile; the crossed circle represents the mean and the inward triangles represent the standard deviation relative to the mean. *** represents statistically significant differences between the quarantine and pre-quarantine conditions as assessed with paired Wilcoxon tests, $p < 0.001$.

Table III. Differences between quarantine and pre-quarantine lifestyle, sleep behavior, procrastination and perceived wellbeing. Statistical significance assessed with paired Wilcoxon test.

| Differential variable | Median | 1 st Q | 3 rd Q | Mean | Standard deviation | Result (p) |
|------------------------------|--------|-------------------|-------------------|-------|--------------------|-------------------|
| DAILY ACTIVITIES (h) | | | | | | |
| Formal study time | 0 | -2 | 0 | -1.01 | 1.57 | Decrease (<0.001) |
| Independent study time | 0 | -2 | 0 | -0.28 | 1.71 | Decrease (0.002) |
| Other daily activities | 2 | 0 | 4 | 1.91 | 2.01 | Increase (<0.001) |
| Phone use time | 1.5 | 0 | 2 | 1.23 | 1.69 | Increase (<0.001) |
| TIME SPENT ONLINE (h) | | | | | | |
| Formal study | 0 | 0 | 1.5 | 0.59 | 1.41 | Increase (<0.001) |
| Independent study | 0 | -0.67 | 0 | -0.04 | 1.21 | No change (0,25) |
| Social media | 0 | 0 | 1.5 | 0.55 | 1.17 | Increase (<0.001) |
| Entertainment | 1 | 0 | 1.5 | 0.81 | 1.27 | Increase (<0.001) |
| SLEEP | | | | | | |
| Sleep time (h) | 0 | 0 | 2 | 0.64 | 1.37 | Increase (<0.001) |
| Going to sleep hour | 0 | 0 | 1 | 0.42 | 1.01 | Increase (<0.001) |
| SQS | -1 | -4 | 2 | -0.61 | 6.26 | Decrease (<0.001) |
| WELLBEING | | | | | | |
| Mood | -1 | -1 | 0 | -0.62 | 1.37 | Decrease (<0.001) |
| Self-org | -1 | -2 | 0 | -0.92 | 1.27 | Decrease (<0.001) |
| Learning efficacy | -1 | -2 | 0 | -0.95 | 1.32 | Decrease (<0.001) |

Differences in sleep perturbing factors

We used the McNemar test to confirm the statistical significance of observed changes in the frequency of sleep-perturbing behaviors during the quarantine compared to before quarantine. We did not observe an increased frequency, but on the contrary, statistically significant decreases for the most common 4 behaviors: variable mealtimes, consumption of coffee and energy drinks before going to sleep, alcohol consumption and tobacco smoking. The results are presented in table IV.

Differences in changes of lifestyle, learning efficacy, procrastination and wellbeing between genders and rural or urban origin

We found statistically significant differences between genders pertaining to the evolution of following indicators: online independent study time, online entertainment time and sleep quality. Namely, while the time spent online for independent study (expressed in hours) decreased for male students [median 0 (-0.92 - 0); mean -0.28 ±1.03], it registered a slight increase for female students [median 0 (-0.67 - 0.67); mean 0.03±1.25], p=0.024. On the other hand, time dedicated to online entertainment increased less among male students [median 0 (0 - 1); mean 0.58±1.16] compared to female students [median 1 (0 - 2); mean 0.87±1.29]. Finally, we observed a more noticeable increase in sleep quality (i.e. higher decrease of SQS) among male students [median -2 (-4 - 0); mean -1.47±5.03] compared to female students [median -1 (-3 - 2); mean -0.38±6.54], p=0.034. Although we failed to prove statistical significance, our results also suggest that

female students had a higher increase in time spent online for formal study [median 0 (0 - 1.5); mean 0.65±1.45] than male students [median 0 (0 - 0.67); mean 0.34±1.23], p=0.062; also, the increase in sleeping time was higher among females [median 0 (0 - 2); mean 0.69±1.41] than males [median 0 (0 - 0); mean 0.44±1.18], p=0.065. It should be noted that pre-quarantine values were similar for students of different genders, except for a significant difference in online formal study (male students studying less than female students).

The only difference regarding students of rural versus urban origin was seen in the change in time spent on social media. As such, there was a higher increase in time spent on social media (expressed in hours) among students of urban origin [median 0.12 (0-1.5); mean 0.6±1.17] compared to students of rural origin [median 0 (0-1); mean 0.31±1.12], p=0.017. There were no statistically significant difference between time spent on social media before quarantine when comparing between students of rural or urban origin.

Students' morale during the quarantine

We have asked the students to rate their (dis) agreement with 8 sentences related to how the quarantine impacted their life. The highest agreement was for the fact that they felt unable to balance leisure with work: 245 students (63.14%) agree or strongly agree. On the other hand, the highest disagreement was towards the idea of wasting less time during quarantine than before: 286 students (73.71%) disagree or strongly disagree. More detailed data are available in table V.

Table IV. Number of students engaging in sleep perturbing factors. Statistical significance assessed with McNemar test.

| Factor | No. before Q | No. after Q | Diff (Q-N) | Gave up | Picked up | P |
|---------------------------------------|--------------|-------------|------------|---------|-----------|--------|
| Variable meal timing | 220 | 193 | -27 | 46 | 19 | 0.001 |
| Coffee or energy drinks 2h before bed | 73 | 40 | -33 | 37 | 4 | <0.001 |
| Alcohol | 40 | 26 | -14 | 23 | 9 | 0.022 |
| Tobacco smoking | 85 | 46 | -39 | 39 | 0 | <0.001 |
| Other substances | 6 | 4 | -2 | 3 | 1 | 0.617 |
| Sedatives | 11 | 5 | -6 | 8 | 2 | 0.114 |
| Other medicine | 26 | 24 | -2 | 3 | 1 | 0.617 |
| Others | 6 | 6 | 0 | 2 | 2 | 1 |

Table V. Students' morale assessment through 8 questions with 5-point Likert scale answers.

| Question | Strongly disagree | Disagree | Neutral | Agree | Strongly agree |
|---|-------------------|----------|---------|-------|----------------|
| Inefficient due to quarantine | 23 | 55 | 98 | 110 | 102 |
| Unable to balance leisure with work | 22 | 61 | 60 | 141 | 104 |
| Unable to self-control | 36 | 78 | 80 | 109 | 85 |
| Desperate due to lack of self-control | 102 | 88 | 79 | 73 | 46 |
| Searched info about escaping Internet addiction | 197 | 96 | 44 | 34 | 17 |
| Quarantine helps me to be more efficient | 94 | 107 | 113 | 55 | 19 |
| I waste less time during the quarantine than before | 172 | 114 | 46 | 36 | 20 |
| The quarantine doesn't affect me | 163 | 99 | 78 | 34 | 14 |

Table VI. Computed correlations between changes in lifestyle, learning efficacy, procrastination and wellbeing. The data use the following format: Spearman correlation coefficient (p-value).

| Differential variable | Mood | Self-org | Learning efficacy | Procrastination |
|--------------------------|-----------------|-----------------|-------------------|-----------------|
| DAILY ACTIVITIES | | | | |
| Formal study time | 0.017 (0.732) | 0.075 (0.140) | 0.069 (0.173) | -0.052 (0.307) |
| Independent study time | 0.139 (0.006) | 0.387 (<0.001) | 0.401 (0) | -0.287 (<0.001) |
| Other daily activities | -0.024 (0.641) | -0.134 (0.008) | -0.101 (0.047) | 0.061 (0.232) |
| Phone use time | -0.188 (<0.001) | -0.295 (<0.001) | -0.253 (<0.001) | 0.231 (<0.001) |
| ONLINE ACTIVITIES | | | | |
| Formal study | -0.043 (0.402) | -0.039 (0.449) | -0.034 (0.500) | -0.017(0.741) |
| Independent study | 0.054 (0.286) | 0.168 (0.001) | 0.162 (0.001) | -0.169 (0.001) |
| Social media | -0.167(0.001) | -0.355 (<0.001) | -0.355 (<0.001) | 0.303 (<0.001) |
| Entertainment | -0.188 (<0.001) | -0.346 (<0.001) | -0.296 (<0.001) | 0.211 (<0.001) |
| SLEEP | | | | |
| Sleep time | -0.017 (0.74) | -0.125 (0.014) | -0.093 (0.066) | 0.071 (0.16) |
| Going to sleep hour | -0.142 (0.005) | -0.188 (<0.001) | -0.199 (<0.001) | 0.117 (0.021) |
| SQS | -0.318 (<0.001) | -0.218 (<0.001) | -0.260 (<0.001) | 0.147 (0.004) |
| WELLBEING | | | | |
| Mood | | 0.537 (<0.001) | 0.523 (<0.001) | -0.281 (<0.001) |
| Self-org | 0.537 (<0.001) | | 0.789 (<0.001) | -0.527 (<0.001) |
| Learning efficacy | 0.523 (<0.001) | 0.789 (<0.001) | | -0.516 (<0.001) |
| Procrastination | -0.281 (<0.001) | -0.527 (<0.001) | -0.516 (<0.001) | |

Correlation between changes in lifestyle, learning efficacy, procrastination, and wellbeing

Finally, we have computed Spearman's correlation coefficient between changes in lifestyle, learning efficacy, procrastination, and wellbeing. Of note are the positive correlations between independent study time and self-organization capacity, and independent study time and learning efficacy (compared to a very weak correlation between formal study time and learning efficacy). Also, change in time spent on social media platforms, time spent on online entertainment and time spent daily on phone are negatively correlated with change in wellbeing indicators across the board. More detailed data are provided in table VI, in the format *correlation coefficient (p-value)*. For all variables, higher values indicate increases during quarantine compared to before the quarantine. Positive correlations towards SQS mean the studied variable was correlated with worse sleep.

Discussion

Using a survey-based study, we have found that the COVID-19 quarantine conditions negatively impacted the lifestyle, learning efficacy and wellbeing of Romanian healthcare students', while also increasing time spent procrastinating. In addition, we have found that the impact of quarantine is influenced by gender and place of origin (urban/rural), and finally we identified negative correlations between wellbeing indicators and procrastination or social media use.

One limitation we encountered stems from the use

of non-probabilistic convenience sampling. This resulted in the over-representation of some demographical categories such as: younger and lower-year students, female students, General Medicine students, and students mainly from two of the 8 analyzed universities. Another limitation is that the questionnaire was self-administered and as such we could not obtain objective data. Moreover, we did not use standardized questionnaires for all the items investigated, except for the PSQI-derived questions, and the questionnaires have not been previously validated.

Stress as a factor explaining changes in wellbeing and procrastination

Numerous studies suggest that, during a quarantine, students have to cope with high levels of stress and a diminished life satisfaction due to quarantine-determined monotony and online classes, in addition to the habitual academic stress [10,22,47–50].

According to some studies, academic procrastination is described as a failure in self-regulation [51], and is strongly correlated with (perceived) stress, anxiety-provoking situations, and life regrets [9,28,52–55]. In our study, we hypothesized that alterations of wellbeing indicators (learning efficacy learning efficacy, mood, self-organization), i.e. negative differences between quarantine and pre-quarantine values, were directly associated with (quarantine v. pre-quarantine) changes in procrastination. We found negative associations between the evolution of procrastination and that of self-organization ($r=-0,52$) and mood ($r=-0,28$), which might indicate that procrastination is linked to reduced life satisfaction, as suggested by

[9,28,55,56]. As expected, changes in mood were rather strongly correlated with the difference between quarantine and pre-quarantine values of self-organization ($r=0.53$) and learning efficacy learning efficacy ($r=0.52$).

Related to our results, Beutel et al. [34], in a study dedicated to procrastination, have reported a positive correlation between procrastination and perceived stress ($r=0.39$) and between procrastination and loneliness ($r=0.27$) and life satisfaction ($r=0.35$). A different study analyzing the psychological effects of COVID-19 quarantines on Chinese school students suggests an upset balance between phone use and physical activity as a possible contributor to the changes observed in our study. Specifically, the authors noticed a nearly fivefold decrease in weekly time dedicated to physical activity time, and that the percentage of sedentary students increased approximately 3 times. Furthermore, screen time during the quarantine increased by about 4 hours per day (1730 minutes per week) [56].

Increased procrastination behaviors are associated with negative consequences ranging from low mood to concerns of mental wellbeing [28,51,53], and subsequently, we hypothesize that increased levels of procrastination are a result of reduced life satisfaction, itself a result of the lockdown [28,34,55].

Yet another possible contributor to a decreased wellbeing of students during the lockdown, and subsequently to procrastination, is mentioned by Twenge et al. [57]. The researchers suggests that 4 hours daily screen time (considered by the authors a moderate level) is associated with lower psychological wellbeing, or more specifically with indicators thereof: inability to finish tasks, lower self-control, less curiosity, less emotional stability [9,20,57].

On a side note, Hong et al. found that dissatisfaction of the need for autonomy is a predictor of proneness to boredom and of increased mobile gaming time, leading, in time, to problematic mobile phone use [58]. Given the fact that, during the quarantine, over 90% of students returned home to live with their families, we hypothesize that decreased independence induced autonomy dissatisfaction that further decreased their wellbeing and caused specific behaviors related to phone and Internet use [28,58].

Lastly, some studies propose that dopamine receptor sensitivity related mechanisms are at play in social media addiction or excessive use [59], due to their short-term provided pleasure [60]. Although we did not initially aim to discuss such physiological mechanisms, the idea that the quarantine-induced increases in time spent online might have had a negative impact on students' wellbeing is a valid future research hypothesis.

Students' morale during quarantine

We assessed the students' morale by asking their (dis)agreement grade with 8 statements. The respondents generally complained of being inefficient, lacking self-control and balance between leisure and work, and

considered that they were affected by quarantine. However, most students also strongly disagreed with the idea of searching information about escaping Internet addiction and being desperate due to lack of self-control, which suggests that they did not perceive a severe increase in Internet use.

Moreover, our study highlighted a decrease in independent (-0.28 h) and formal (-1.01 h) study time, an increase in phone use (0.55 h), and online entertainment (0.81 h). Additionally, time spent with other daily activities increased by almost 2 hours.

Similar to our results, a survey of Saudi medical students [21] has found deterioration of work performance, lack of concentration during studying and a decrease in overall study time, a part of the students also reporting difficulty in mental calculations.

Students' sleep quality during quarantine

Unlike most studies on poor sleep quality in quarantine [61,62], we observed a modest, but statistically significant, improvement in sleep quality (mean decrease of SQS score=-0,61, consistent with a better sleep quality), and an increase in sleep duration (mean increase=0,64 h). However, students tended to go to sleep later during quarantine (on average 25 minutes later), which is similar to the results of previous studies [62,63]. Those changes might be due to most of the students (352 students, 90,72%) returning home and having online courses, as such the time spent daily for going to and coming from University courses has been nullified, and the need to wake up early (and incidentally, to go to sleep early) disappeared. However, increasing free time during quarantine might explain the improvement in sleep quality and duration. Otherwise, students were able to skip hours online, which could result in better sleep quality and duration.

The statistically significant improvement in sleep quality is nonetheless unexpected. While the negative aspects of quarantine (boredom, frustration, lack of social support, fear of uncertainty) are likely to have contributed to the previously mentioned decrease in wellbeing, they seem to have had a minimal effect on sleep quality, possibly outweighed by the comfort of sleeping at home.

Differences between students or rural and urban origin, pertaining to social media use

The only statistically significant difference between students of rural or urban origin we observed was between the increases in time spent on social media (namely, students of rural origin spent on average 18 minutes more on social media during the quarantine, while students of urban origin spent the double of that, 36 minutes). Together with the fact that most students went home due to quarantine, we further hypothesize that this difference comes from the fact that in rural areas, physical activity is more accessible than in urban areas and as such, students of rural origin (which also lived in rural areas during the quarantine) went outside more and thus spent less time on social media [64,65].

Conclusion

The long-term quarantine imposed in response to the COVID-19 pandemic determined lifestyle, learning efficacy and wellbeing changes of Romanian healthcare students, which previous studies linked to heightened levels of stress, monotony, boredom and frustration. Specifically, the lockdown caused an increased phone and social media use, at the expense of formal and independent study, as well as deteriorations in mood, self-organization capacity and learning efficacy learning efficacy, and increased procrastination behaviors. However, the increase in social media use was slightly lower for students of rural origin.

Generally, students complained of being inefficient, lacking self-control and having difficulty in balancing work and leisure. On the other hand, the quarantine's effects on sleep quality and duration were contrary to those of other studies, the two parameters registering a moderate improvement.

In conclusion, our study draws attention to the problem of quarantine-induced deteriorations of wellbeing and learning capacity in a category of students who are already considered to experience increased levels of study-related stress. Subsequently, we propose social and mental health support as a means of minimizing the negative impact of quarantine.

References

- Lin X, Gong Z, Xiao Z, Xiong J, Fan B, Liu J. Novel Coronavirus Pneumonia Outbreak in 2019: Computed Tomographic Findings in Two Cases. *Korean J Radiol.* 2020;21:365–368.
- Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. *J Adv Res.* 2020;24:91–98.
- World Health Organization. Pneumonia of Unknown Cause – China. Available online: <https://www.who.int/csr/don/05-january-2020-pneumonia-of-unknown-cause-china/en/>.
- Sun P, Lu X, Xu C, Sun W, Pan B. Understanding of COVID-19 based on current evidence. *J Med Virol.* 2020;92:548–551.
- Centrul Național de Supraveghere și Control al Bolilor Transmisibile Analiza Cazuri Confirmate (261). Pana La 18 03 2020 [National Center of Control of Communicable Diseases Analysis of Confirmed Cases] [Romanian] Available online: <https://www.cnscbt.ro/index.php/analiza-cazuri-confirmate-covid19/1539-analiza-cazuri-confirmate-261-pana-la-18-03-2020> (accessed on 6 June 2022).
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet.* 2020;395:912–920.
- Garfin DR, Silver RC, Holman EA. The novel coronavirus (COVID-2019) outbreak: Amplification of public health consequences by media exposure. *Health Psychol.* 2020;39:355–357.
- Shi L, Lu ZA, Que JY, Huang XL, Liu L, Ran MS, et al. Prevalence of and Risk Factors Associated with Mental Health Symptoms Among the General Population in China During the Coronavirus Disease 2019 Pandemic. *JAMA Netw Open.* 2020;3:e2014053.
- Sifat RI, Ruponty MM, Rahim Shuvo MK, Chowdhury M, Suha SM. Impact of COVID-19 pandemic on the mental health of school-going adolescents: insights from Dhaka City, Bangladesh. *Heliyon.* 2022;8:e09223.
- Yan L, Gan Y, Ding X, Wu J, Duan H. The relationship between perceived stress and emotional distress during the COVID-19 outbreak: Effects of boredom proneness and coping style. *J Anxiety Disord.* 2021;77:102328.
- Blendon RJ, Benson JM, DesRoches CM, Raleigh E, Taylor-Clark K. The public's response to severe acute respiratory syndrome in Toronto and the United States. *Clin Infect Dis.* 2004;38:925–931.
- DiGiovanni C, Conley J, Chiu D, Zaborski J. Factors influencing compliance with quarantine in Toronto during the 2003 SARS outbreak. *Biosecur Bioterror.* 2004;2:265–272.
- Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS control and psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis.* 2004;10:1206–1212.
- Lee S, Chan LY, Chau AM, Kwok KP, Kleinman A. The experience of SARS-related stigma at Amoy Gardens. *Soc Sci Med.* 2005;61:2038–2046.
- Bai Y, Lin CC, Lin CY, Chen JY, Chue CM, Chou P. Survey of stress reactions among health care workers involved with the SARS outbreak. *Psychiatr Serv.* 2004;55:1055–1057.
- Maunder R, Hunter J, Vincent L, Bennett J, Peladeau N, Leszcz M, et al. The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital. *CMAJ.* 2003;168:1245–1251.
- Desclaux A, Badji D, Ndione AG, Sow K. Accepted monitoring or endured quarantine? Ebola contacts' perceptions in Senegal. *Soc Sci Med.* 2017;178:38–45.
- Jose R, Holman EA, Silver RC. The Importance of the neighborhood in the 2014 Ebola outbreak in the United States: Distress, worry, and functioning. *Health Psychol.* 2017;36:1181–1185.
- Korczak DJ, Madigan S, Colasanto M. Children's Physical Activity and Depression: A Meta-analysis. *Pediatrics.* 2017;139:e20162266.
- Ullah I, Islam MS, Ali S, Jamil H, Tahir MJ, Arsh A, et al. Insufficient Physical Activity and Sedentary Behaviors among Medical Students during the COVID-19 Lockdown: Findings from a Cross-Sectional Study in Pakistan. *Int J Environ Res Public Health.* 2021;18:10257.
- Meo SA, Abukhalaf AA, Alomar AA, Sattar K, Klonoff DC. COVID-19 Pandemic: Impact of Quarantine on Medical Students' Mental Wellbeing and Learning Behaviors. *Pak J Med Sci.* 2020;36(COVID19-S4):S43–S48.
- Dumitrache L, Stănculescu E, Nae M, Dumbrăveanu D, Simion G, Taloș AM, et al. Post-Lockdown Effects on Students' Mental Health in Romania: Perceived Stress,

- Missing Daily Social Interactions, and Boredom Proneness. *Int J Environ Res Public Health*. 2021;18:8599.
23. Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on College Students' Mental Health in the United States: Interview Survey Study. *J Med Internet Res*. 2020;22:e21279.
 24. Satici B, Saricali M, Satici SA, Griffiths MD. Intolerance of Uncertainty and Mental Wellbeing: Serial Mediation by Rumination and Fear of COVID-19. *Int J Ment Health Addict*. 2020;1–12.
 25. Schimmenti A, Billieux J, Starcevic V. The Four Horsemen of Fear: An Integrated Model of Understanding Fear Experiences During the Covid-19 Pandemic. *Clin Neuropsychiatry*. 2020;17:41–45.
 26. Hall G, Laddu DR, Phillips SA, Lavie CJ, Arena R. A tale of two pandemics: how will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Prog Cardiovasc Dis*. 2021;64:108–110.
 27. Gröpel P, Steel P. A mega-trial investigation of goal setting, interest enhancement, and energy on procrastination. *Personality and Individual Differences*. 2008;45:406–411.
 28. Wang Q, Kou Z, Du Y, Wang K, Xu Y. Academic Procrastination and Negative Emotions Among Adolescents During the COVID-19 Pandemic: The Mediating and Buffering Effects of Online-Shopping Addiction. *Front Psychol*. 2022;12:789505.
 29. Azad MC, Fraser K, Rumana N, Abdullah AF, Shahana N, Hanly PJ, et al. Sleep disturbances among medical students: a global perspective. *J Clin Sleep Med*. 2015;11:69–74.
 30. Meo SA, Abukhalaf DAA, Alomar AA, Sattar K, Klonoff DC. Burnout syndrome and associated factors among medical students: a cross-sectional study. *Clin (Sao Paulo)*. 2012;67:573–580.
 31. Mohd Sidik S, Rampal L, Kaneson N. Prevalence of emotional disorders among medical students in a Malaysian university. *Asia Pac Fam Med*. 2003;2:213–217.
 32. Zeng W, Chen R, Wang X, Zhang Q, Deng W. Prevalence of mental health problems among medical students in China. *Medicine (Baltimore)*. 2019;98:e15337.
 33. Xiao H, Zhang Y, Kong D, Li S, Yang N. The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit*. 2020;26:e923549.
 34. Beutel ME, Klein EM, Aufenanger S, Brähler E, Dreier M, Müller KW, et al. Procrastination, Distress and Life Satisfaction across the Age Range – A German Representative Community Study. *PLoS One*. 2016;11:e0148054.
 35. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer D.J. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28:193–213.
 36. Katagiri R, Asakura K, Kobayashi S, Suga H, Sasaki S. Low intake of vegetables, high intake of confectionary, and unhealthy eating habits are associated with poor sleep quality among middle-aged female Japanese workers. *J Occup Health*. 2014;56:359–368.
 37. St-Onge MP, Mikic A, Pietrolungo CE. Effects of Diet on Sleep Quality. *Adv Nutr*. 2016;7:938–949.
 38. Lohsoonthorn V, Khidir H, Casillas G, Lertmaharit S, Tadesse MG, Pensuksan WC, et al. Sleep quality and sleep patterns in relation to consumption of energy drinks, caffeinated beverages, and other stimulants among Thai college students. *Sleep Breath*. 2013;17:1017–1028.
 39. Mwape RK, Mulenga D. Consumption of Energy Drinks and Their Effects on Sleep Quality among Students at the Copperbelt University School of Medicine in Zambia. *Sleep Disord*. 2019;2019:3434507.
 40. Park SY, Oh MK, Lee BS, Kim HG, Lee WJ, Lee JH, et al. The Effects of Alcohol on Quality of Sleep. *Korean J Fam Med*. 2015;36:294–299.
 41. Roehrs T, Roth T. Sleep, sleepiness, sleep disorders and alcohol use and abuse. *Sleep Med Rev*. 2001;5:287–297.
 42. Liao Y, Xie L, Chen X, Kelly BC, Qi C, Pan C, et al. Sleep quality in cigarette smokers and nonsmokers: findings from the general population in central China. *BMC Public Health*. 2019;19:808.
 43. Wetter DW, Young TB, Bidwell TR, Badr MS, Palta M. Smoking as a risk factor for sleep-disordered breathing. *Arch Intern Med*. 1994;154:2219–2224.
 44. Zhabenko O, Austic E, Conroy DA, Ehrlich P, Singh V, Epstein-Ngo Q, et al. Substance Use as a Risk Factor for Sleep Problems among Adolescents Presenting to the Emergency Department. *J Addict Med*. 2016;10:331–338.
 45. Min Y, Kirkwood CK, Mays DP, Slattum, PW. The Effect of Sleep Medication Use and Poor Sleep Quality on Risk of Falls in Community-Dwelling Older Adults in the US: A Prospective Cohort Study. *Drugs Aging*. 2016;33:151–158.
 46. R Core Team R: A Language and Environment for Statistical Computing; R Foundation for Statistical Computing, Vienna, Austria, 2021.
 47. Al-Rabiaah A, Temsah MH, Al-Eyadhy AA, Hasan GM, Al-Zamil F, Al-Subaie S, et al. Middle East Respiratory Syndrome-Corona Virus (MERS-CoV) associated stress among medical students at a university teaching hospital in Saudi Arabia. *J Infect Public Health*. 2020;13:687–691.
 48. Odriozola-González P, Planchuelo-Gómez Á, Irujo MJ, de Luis-García R. Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. *Psychiatry Res*. 2020;290:113108.
 49. Sahu P. Closure of Universities Due to Coronavirus Disease 2019 (COVID-19): Impact on Education and Mental Health of Students and Academic Staff. *Cureus* 2020;12:e7541.
 50. Zhai Y, Du X. Mental health care for international Chinese students affected by the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7:e22.
 51. Steel P. The nature of procrastination: a meta-analytic and theoretical review of quintessential self-regulatory failure. *Psychol Bull*. 2007;133:65–94.
 52. Farris SG, Kibbey MM, Fedorenko EJ, DiBello AM. A Qualitative Study of COVID-19 Distress in University Students. *Emerg Adulthood* 2021;9:462–478.
 53. Ferrari JR, Barnes KL, Steel P. Life Regrets by Avoidant and Arousal Procrastinators: Why Put off Today What You Will Regret Tomorrow? *J Individ Differ*. 2009;30:163–168.

54. Hen M, Goroshit M. General and Life-Domain Procrastination in Highly Educated Adults in Israel. *Front Psychol.* 2018;9:1173.
55. Rebetz MM, Rochat L, Gay P, Van der Linden M. Validation of a French version of the pure procrastination scale (PPS). *Compr Psychiatry.* 2014;55:1442–1447.
56. Xiang M, Zhang Z, Kuwahara K. Impact of COVID-19 pandemic on children and adolescents' lifestyle behavior larger than expected. *Prog Cardiovasc Dis.* 2020;63:531–532.
57. Twenge JM, Campbell WK. Associations between screen time and lower psychological well-being among children and adolescents: Evidence from a population-based study. *Prev Med Rep.* 2018;12:271–283.
58. Hong W, Liu RD, Ding Y, Zhen R, Jiang R, Fu X. Autonomy Need Dissatisfaction in Daily Life and Problematic Mobile Phone Use: The Mediating Roles of Boredom Proneness and Mobile Phone Gaming. *Int J Environ Res Public Health.* 2020;17:5305.
59. Guedes E, Sancassiani F, Carta MG, Campos C, Machado S, King AL, et al. Internet Addiction and Excessive Social Networks Use: What About Facebook? *Clin Pract Epidemiol Ment Health.* 2016;12:43–48.
60. Christensen JF. Pleasure junkies all around! why it matters and why 'the arts' might be the answer: a biopsychological perspective. *Proc Biol Sci.* 2017;284:20162837.
61. Fu W, Wang C, Zou L, Guo Y, Lu Z, Yan S, et al. Psychological health, sleep quality, and coping styles to stress facing the COVID-19 in Wuhan, China. *Transl Psychiatry.* 2020;10:225.
62. Marelli S, Castelnovo A, Somma A, Castronovo V, Mombelli S, Bottoni D, et al. Impact of COVID-19 lockdown on sleep quality in university students and administration staff. *J Neurol.* 2021;268:8–15.
63. Cellini N, Canale N, Mioni G, Costa S. Changes in sleep pattern, sense of time and digital media use during COVID-19 lockdown in Italy. *J Sleep Res.* 2020;29:e13074.
64. Peen J, Dekker J, Schoevers RA, Have MT, de Graaf R, Beekman AT. Is the Prevalence of psychiatric disorders associated with urbanization? *Soc Psychiatry Psychiatr Epidemiol.* 2007;42:984–989.
65. Stavropoulos V, Alexandraki K, Motti-Stefanidi F. Recognizing internet addiction: prevalence and relationship to academic achievement in adolescents enrolled in urban and rural Greek high schools. *J Adolesc.* 2013;36:565–576.